Images in Cardiovascular Disease

Loeffler Endocarditis: A Diagnosis Made with Cardiovascular Magnetic Resonance

Dinesh K Kalra , MD, Jeffrey Park , MD, Mohamad Hemu , MD, and Alan Goldberg , MD

Department of Internal Medicine, Rush University Medical Center, Chicago, IL, USA

OPEN ACCESS

Received: Nov 5, 2018

Revised: Dec 19, 2018

Accepted: Dec 26, 2018

Address for Correspondence:

Dinesh K Kalra, MD

Division of Cardiology, Department of Internal Medicine, Rush University Medical Center, 1717 W Congress Parkway, Kellogg Suite 320, Chicago, IL 60612, USA.
E-mail: dinesh_kalra@rush.edu

Copyright © 2019 Korean Society of Echocardiography

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Dinesh K Kalra

<https://orcid.org/0000-0001-5254-4067>

Jeffrey Park

<https://orcid.org/0000-0002-2117-7118>

Mohamad Hemu

<https://orcid.org/0000-0003-3059-7437>

Alan Goldberg

<https://orcid.org/0000-0003-2257-119X>

Conflict of Interest

The authors have no financial conflicts of interest.

A 61-year-old Caucasian man with recently diagnosed eosinophilic chronic myelomonocytic leukemia presented with fevers and right-sided heart failure that failed to respond to diuretics. He denied any history of travel to the tropics. Exam was remarkable for pallor, edema and S4 gallop. He had leukocytosis ($54 \times 10^9/L$; 55% eosinophilic predominance), anemia and thrombocytopenia ($3 \times 10^9/L$). Echocardiogram showed grade 2 diastolic dysfunction, moderate mitral regurgitation and elevated right heart pressure. There was apical left ventricular thickening concerning for tumor, laminated thrombus or focal hypertrophy (**Figure 1, Movie 1**). Given severe thrombocytopenia, endomyocardial biopsy and anticoagulation were contraindicated. Cardiovascular magnetic resonance (CMR, **Figure 2, Movie 2, 3**) showed increased focal apical wall thickness of 22 mm with hypokinesis of this segment (**Figure 2A**). First pass perfusion revealed hypoperfused apical and lateral endocardium with a layer of superimposed clot (**Figure 2B**). Late gadolinium enhancement images showed hyperintense apical fibrosis extending from the endocardium to mid myocardium, covered by a cap where laminated clot was present producing a “double V” sign (**Figure 2C**). This was also confirmed on the 3-chamber early gadolinium enhancement image with long TI of 600 ms that showed apical thrombus distinct from the underlying myocardium (**Figure 2D**).

Figure 1. Echo (apical 4 chamber) with perflutren contrast showing a 2.4×3.8 cm non-perfused mass at the LV apex suggestive of thrombus or infiltrative tumor. LA: left atrium, LV: left ventricle, RV: right ventricle.

Figure 2. (A) Cardiovascular magnetic resonance – Steady State Free Precession (SSFP) still image (apical 2 chamber) showing normal chamber sizes with increased focal apical wall thickness of 22 mm and hypokinesis of this segment. A small pericardial effusion is also seen. (B) First pass perfusion shows hypoperfused apical and lateral endocardium along with a layer of clot lining these segments. (C) Three-chamber late gadolinium enhancement shows fibrosis in the apex in the endocardium extending to the mid myocardium in the shape of a cap. The laminated hypointense clot (arrowhead) produces a “double V” sign when contrasted with the underlying hyperintense fibrosis (asterisk). (D) Three-chamber early gadolinium enhancement with long TI of 600 ms shows dark hypointense thrombus at the apex distinct from the underlying scarred myocardium. LA: left atrium, LV: left ventricle, RV: right ventricle.

Loeffler endocarditis is a rare type of restrictive cardiomyopathy that follows a two-stage process.^{1,2)} Early manifestations include symptoms of fevers, dyspnea and swelling. During this phase, infiltration of myocardium and release of eosinophilic major basic protein and reactive oxygen species promote endothelial and myocyte injury.³⁾ The later stages are marked by myocardial fibrosis and thrombus formation causing a restrictive cardiomyopathy.^{4,5)} Our patient was started on imatinib and steroids and his symptoms improved over 2 weeks. Unfortunately, he died from bacterial sepsis after that.

SUPPLEMENTARY MATERIALS

Movie 1

Apical 4-chamber echo showing laminated mass at left ventricular apex.

[Click here to view](#)

Movie 2

Cardiovascular magnetic resonance – 2-chamber Steady State Free Precession cine imaging showing Focal thickening at the left ventricular apex (22 mm) along with hypokinesis.

[Click here to view](#)

Movie 3

Cardiovascular magnetic resonance – Apical short-axis first pass perfusion imaging demonstrating hypoperfusion of the subendocardium and a layer of overlying thrombus.

[Click here to view](#)

REFERENCES

1. Mankad R, Bonnicksen C, Mankad S. Hypereosinophilic syndrome: cardiac diagnosis and management. *Heart* 2016;102:100-6.
[PUBMED](#) | [CROSSREF](#)

2. Ogbogu PU, Rosing DR, Horne MK 3rd. Cardiovascular manifestations of hypereosinophilic syndromes. *Immunol Allergy Clin North Am* 2007;27:457-75.
[PUBMED](#) | [CROSSREF](#)
3. Spry CJ, Tai PC, Davies J. The cardiotoxicity of eosinophils. *Postgrad Med J* 1983;59:147-53.
[PUBMED](#) | [CROSSREF](#)
4. Beedupalli J, Modi K. Early-stage Loeffler's endocarditis with isolated right ventricular involvement: management, long-term follow-up, and review of literature. *Echocardiography* 2016;33:1422-7.
[PUBMED](#) | [CROSSREF](#)
5. Kleinfeldt T, Ince H, Nienaber CA. Hypereosinophilic syndrome: a rare case of Loeffler's endocarditis documented in cardiac MRI. *Int J Cardiol* 2011;149:e30-2.
[PUBMED](#) | [CROSSREF](#)